



Entrez PubMed

Nucleotide

Protein

Genome

Structure

PMC

Journals

Bo

Search PubMed



for



Clear

☒ Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract



Show:

20



Sort



Send to



Text

Text Version

☐ 1: Surgery. 1993 Jun; 113(6): 676-82.

Related Articles, Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

Inhibition of tumor cell adhesion to lymph nodes by laminin-related peptide and neuraminidase.

Islam SM, Whalen GF, Sharif SF.

Department of Surgery, Cornell Medical College, New York, N.Y.

BACKGROUND. Adhesion to lymph nodes, rather than growth stimulation, accounted for preferential colonization of lymph nodes by a metastatic B16 melanoma. We investigated these adhesive interactions. **METHODS.** Four classes of molecules were tested for inhibition of melanoma adhesion to cryostat sections of lymph node. **RESULTS.** Calcium chelators ethylenediaminetetraacetic acid and ethyleneglycol-bis-(beta-aminoethylether)-N,N,N',N'-tetra acetic acid completely inhibited adhesion (50% adhesion, half-maximal inhibition, at 1 to 3 mmol/L). Cytochalasin B, which impairs contractile microfilaments, inhibited adhesion (60% adhesion at .001 mmol/L, 28% at .01 mmol/L). Colchicine, which disaggregates microtubules, had a similar effect (20% at .01 mmol/L, lowest dose tested). Trypsin slightly increased adhesion (125% adhesion at 10 micrograms/ml). Neuraminidase, which removed sialic acid residues, inhibited it (50% adhesion at 5 micrograms/ml). Gly-arg-gly-asp-ser, a peptide with a cell binding sequence of fibronectin, did not consistently inhibit adhesion (69% adhesion at 0.1 mg/ml, 83% adhesion at 1 mg/ml) or substantially differ from gly-arg-gly-glu-ser-pro (59% adhesion at 0.1 mg/ml, 90% adhesion at 1 mg/ml). In contrast, a peptide with a cell binding region of laminin (tyr-ile-gly-ser-arg) inhibited adhesion (50% adhesion at .05 mg/ml). **CONCLUSIONS.** Tumor cell-lymph node adhesion is a calcium-dependent process, requiring a functional cytoskeleton, that is mediated by both sialic acid moieties and trypsin-resistant, laminin-related, adhesion molecules.

PMID: 8506527 [PubMed - indexed for MEDLINE]

Display

Abstract



Show:

20



Sort



Send to



Text

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)